

**A PARTIAL LEUCISM CASE IN *COLUMBINA PICUI* (TEMMINCK 1813)
(BIRDS: COLUMBIIFORMS), IN SOUTH OF BRAZIL**

*Luiz Liberato Costa Corrêa^{1,2};
Dariane Evangelho Silva^{1,2}
Stefan Vilges de Oliveira^{2,3}

RESUMO

Apresentamos o registro de um caso de leucismo parcial em *Columbina picui* no município de São Sepé, Rio Grande do Sul, Brasil.

Palavras chave: Anomalia genética, Leucismo, Rolinha-picuí, São Sepé, Rio Grande do Sul.

UM CASO DE LEUCISMO PARCIAL EM *COLUMBINA PICUI* (Temminck 1813) (Aves: Columbiformes), NO SUL DO BRASIL

ABSTRACT

The record of partial leucism in *Columbina picui* is presented from the municipality of São Sepé, Rio Grande do Sul, Brazil.

Keywords: Genetic disorder, Leucism , Picui Ground-Dove, São Sepé, Rio Grande do Sul.

¹ PPG em Ambiente e Desenvolvimento, Centro Universitário UNIVATES, Avenida Avelino Tallini, 171, CEP 95900-000, Lajeado – RS, Brasil.

² Organização Não Governamental – Interação de Trabalhos Ambientais Caçapava do Sul – RS, Brasil.

³ PPG em Medicina Tropical, Epidemiologia das Doenças Infecciosas e Parasitárias Núcleo de Medicina Tropical, Universidade de Brasília, Brasília – DF, Brasil.

*Autor de contato: lc_correa@yahoo.com.br

INTRODUCTION

Birds can present a wide range of color variation, some from biologic/ecologic character, as for example the sexual dimorphism which is showed through chromatic variation between male and female. Even different color patterns can be seen in young individuals and adults (BUCKLEY, 1982), as well individual variation occurring due to genetic mutations as xantocroism, melanism, cyanism, erytrism, leucism and albinism (SIGRIST, 2006; URCULA, 2011). The partial or total pigment loss in feathers of birds is mistakenly given as albinism by some authors, however this occurs only in the total absence of melanin. In other words not only in feathers, occurring also in the eyes and skin due to lack of tyrosinase enzyme, caused by genetic mutation (GROSS, 1965; GRILLI *et al.*, 2006; VAN GROUW, 2006).

The leucism, study object of this article, is caused by the partial or total lack of eumelanin and feomelanin in the birds feathers. Many times it is restricted to some parts of the plumage, as it havetyros in and presents dark iris, not pink or reddish as occurs in albino individuals (GROSS, 1965, AJALA and MIKKOLA, 1997; VAN GROUW, 2006; CARBÓ-RAMÍREZ *et al.*, 2011; URCULA, 2011).

Several leucism registers have been done on birds: *Thalassarche melanophris* e *Daption capense* (MANCINI *et al.*, 2010), *Passer domesticus* (CORRÊA *et al.*, 2011), *Volatinia jacarina* (GAIOTTI *et al.*, 2011), *Paroaria coronata* (CORRÊA *et al.*, 2012), and *Pygochelidon cyanoleuca* (GODOY, 2012), and these aim to remark the variation in color pattern and behavior. In Columbiforms, some chromatic mutations are reported by Mallet-Rodrigues (1995) in *Columbina talpacoti*, González-Acuña (2004) *Zenaida auriculata*, SANTOS *et al.* (2011) *Patagioenas picazuro* and Urcula (2011) *Columba maculosa*, *Z. auriculata* and one albino specimen in *Columbina picui*. However, *C. picui* leucism has not been reported by literature in the Southern region of Brazil.

Columbina picui commonly known as Picui Ground-dove is a Columbidae family small body size bird, which occurs from Colombia to Bolivia, Chile, Argentina, Uruguay and Brazil (SICK, 1997). This bird is known by its remarkable white tail feathers and by the contrast of black and white in the wing feathers, grayish color patterns among feathers and plumes, dark red tarsi and black beak (BELTON, 1994; SICK, 1997). In Rio Grande do Sul is considered widespread in rural and urban zones, commonly found in pairs or small bunches (BELTON, 1994).

This note presents a partial leucism case in *Columbina picui* in urban area of the municipality of São Sepé, central region of the State of Rio Grande do Sul, Brazil.

OCCURRENCE DESCRIPTION

One mutant individual of *C. picui* was seen by January 24th, 2013, in the morning (around 10 am) in a urbanized area (S 30° 09', 05" W 53° 34', 00"S), in the municipality of São Sepé, central region of Rio Grande do Sul. At the moment of this observation, this specimen was foraging in the sidewalk at the Kurtz District. The specimen was besides another individual of *C. picui*, with normal coloration (Figure 1).

For the data collection about this specimen, the effort applied was about 6 hours of direct observation during the same day of the first register using binoculars (10 x 50 lenses). Notes were taken about the general living and behavior of this bird.

The partially leucistic *C. picui* specimen is an adult with no identified gender. It shows depigmentation in part of the wing feathers, some remiges and coverts, also in some plumes and feathers by the chest and back (Figure 2). These feathers and plumes are grayish in normal individuals (BELTON, 1994). Eyes, iris, tarsi and beak, still in normal pigmentation according to Belton (1994) and Sick (1997) descriptions. By the time of observation this *C. picui* specimen didn't showed any abnormal behavior in relation to the other *C. picui* specimens.



Figure 1: Partially leucistic *C. picui* specimen (right), accompanied by one specimen in standard colors (left), in urban area of São Sepé municipality, RS. Photo: Luiz Corrêa.



Figure 2: Partially leucistic *C. picui* specimen in urban area of São Sepé, RS. Can be seen depigmentation in some feathers, plumes, coverts and remiges. Photo: Luiz Corrêa.

Is remarkable the fact that in the same surroundings (around 600 meters) was reported a leucistic specimen of *Passer domesticus*, by Corrêa *et al.* (2010). The occurrence of leucism seems to have strong relation with the urban environment (CORRÊA *et al.*, 2010; RIBEIRO and GOGLIATH, 2012). This probably can be explained by the low abundance of predators (ELLEGREN *et al.*, 1997), fact that makes natural selection less intense in urban centers, which can justify the more leucism occurrence in these areas. Some authors give more attention in the potential concentration of mutagenic compounds in these environments (MØLLER and MOUSSEAU, 2001). As this is the second report in this area, we suggest additional investigation on these constraints.

Finishing, is important to keep monitoring this free living mutant specimen by mid and long term, collecting relevant information that can explain these occurrence patterns, identifying potential causes and effects.

REFERENCES

ALAJA, P.; MIKKOLA, H. Albinism in the Great Gray Owl (*Strix nebulosa*) and other owls, (In): DUNCAN, J. R.; JOHNSON, D. H. ; NICHOLLS T. H. (eds.) *Biology and conservation of owls of the Northern hemisphere*. Washington, D.C.: USDA Forest Service (General Technical Report NC-190), Pp. 33-37.1997.

BELTON, W. **Aves do Rio Grande do Sul: distribuição e biologia.** São Leopoldo: Editora Unisinos, 1994. 584p

BUCKLEY, P. A. Avian genetics. (In): PETRAK, M. L. (Ed.). **Diseases of cage and aviary birds**. Lea and Febiger, Philadelphia, USA, Pp. 21-110. 1982.

CARBÓ-RAMÍREZ, P.; ROMERO-GONZÁLEZ, P.; ZURIA, I. Primer reporte para México de coloración aberrante (*leucismo parcial*) en el cuitlaco-che pico curvo (*Toxostoma curvirostre*). **Huitzil**, v. 12, n. 1, p.1-4, 2011.

CORRÊA, L. L. C.; SILVA, D. E. ; TRINDADE, A. de O. ; OLIVEIRA, S. V. de. Registro de Leucismo em Pardal (*Passer domesticus*) (Lineaeus, 1758), para o Sul do Brasil. **Biodiversidade Pampeana**, v. 9, n.1, p.12-15, 2011.

CORRÊA, L. L. C.; SILVA, D. E. ; FERLA, N. J. ; SEIXAS, A. L. da R.; OLIVEIRA, S. V. de. Registro de leucismo em Cardeal *Paroaria coronata* (Miller, 1776) no Sul do Brasil. **Revista de Ciências Ambientais**, v. 6, n. 2, p. 73- 79, 2012.

ELLEGREN, H.; LINDGREN, G.; PRIMMER, C. R.; MØLLER A. P. Fitness lossand germline mutations in Barn Swallows breeding in Chernobyl. **Nature**, v. 389, n. 9, p. 593-596, 1997.

GAIOTTI, M. G., PINHO, J. B.; GRANGEIRO, D. New record of aberrant plumage in Blue-black Grassquit (*Volatinia jacarina* Linnaeus, 1766, aves: Emberizidae). **Brazilian Journal of Biology**, v. 71, n. 2, p. 567, 2011.

GODOY, F. I. D. Registro de leucismo total em andorinha-pequena-de-casa (*Pygochelidon cyanoleuca*). **Atualidades Ornitológicas**, n. 168, p.18-19, 2012.

GONZÁLEZ-ACUÑA, D. Albinismo en un ejemplar de *Zenaida auriculata* (Des Murs, 1847) en Ñuble. **Boletín Chileno de Ornitología**, n. 10, p. 25-26, 2004.

GRILLI, P. G.; MOSCHIONE, F. N.; BURGOS, F. G. Leucismo parcial en pepitero de collar *Saltator aurantiirostris* en Santa Bárbara, Jujuy, Argentina. **Cotinga**, n. 25, p. 89 90, 2006.

GROSS, A. O. The incidence of albinism in North American birds. **Bird-Banding**, v. 36, n. 2, p. 67-71, 1965.

GROUW, H. VAN. Not every white bird is an albino: sense and nonsense about colour aberrations in birds. **Dutch Birding**, n. 28, p. 79-89, 2006.

MALLET-RODRIGUES, F. Um registro de albinismo em *Columbina talpacoti* (Temincki, 1811) para o Rio de Janeiro. **Atualidades Ornitológicas**, n. 64, p. 12, 1995.

MANCINI, L. P.; JIMÉNEZ, S.; NEVES, T.; BUGONI, L. Records of leucism in albatrosses and petrels (Procellariiformes) in the South Atlantic Ocean. **Revista Brasileira de Ornitologia**, v. 18, n. 3, p. 245-248, 2010.

MØLLER, A. P. ; MOUSSEAU, T. A. Albinism and phenotype of Barn Swallows (*Hirundo rustica*) from Chernobyl. **Evolution**, v. 55, n. 10, p. 2097-2104, 2001.

URCOLA, M. R. Aberraciones cromáticas en aves de la colección ornitológica Del Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”. **Revista del Museo Argentino Ciencias Naturales**, v. 13, n.2, p. 221-228, 2011.

RIBEIRO, L. D. B.; GOGLIAT, H. M. Um caso de leucismo em pardal, *Passer domesticus* (Linnaeus, 1758) em uma ilha do rio São Francisco, nordeste do Brasil. **Biotemas**, v. 25, n.1, p. 187-190, 2012.

SANTOS, K. K. , LOMBARDI, V. T.; D'ANGELO NETO, S.; MIGUEL, M.; FAETTI, R. G. Registros de plumagens aberrantes em *Patagioenas picazuro* (Columbiformes, Columbidae), *Knipolegus lophotes* (Passeriformes, Tyrannidae) e *Turdus rufiventris* (Passeriformes, Turdidae) no estado de Minas Gerais. **Atualidades Ornitológicas**, n.160, p. 24-25, 2011.

SICK, H. **Ornitologia brasileira**. Rio de Janeiro: Nova Fronteira. 1997. 862p.

SIGRIST, T. **Aves do Brasil, uma visão artística**. São Paulo: Fosfértil. 2006. 672p.