

## Knowledge and adherence to preventive measures for ventilator-associated pneumonia during the COVID-19 pandemic

*Conhecimento e adesão às medidas preventivas da pneumonia associada à ventilação mecânica na pandemia de COVID-19*

*Conocimiento y adherencia a las medidas preventivas de la neumonía asociada a ventilador en la pandemia de COVID-19*

<https://doi.org/10.17058/reci.v14i4.19180>

Received: 02/21/2024



Accepted: 10/17/2024

Available online: 12/18/2024

**Corresponding Author:**

Hélio Galdino Júnior  
helio\_junior@ufg.br

Address: Rua 1, Chácaras Alto da Glória,  
Goiânia, Goiás, Brasil.

Ana Clara Alves Campos<sup>1</sup> ;  
Jessica de Oliveira Montebello<sup>1</sup> ;  
George Oliveira Silva<sup>1</sup> ;  
Hélio Galdino Júnior<sup>1</sup> 

<sup>1</sup> Universidade Federal de Goiás, Goiânia, GO, Brasil.

### ABSTRACT

**Background and Objectives:** adherence to measures to prevent mechanical ventilation-associated pneumonia (VAP) significantly reduces this hospital infection, and knowledge of the measures is essential to ensure adherence. COVID-19 has increased the demand for intensive care and increases the risks and severity of VAP. Thus, the objective was to assess the knowledge referred to about the VAP prevention bundle and its associated factors and multidisciplinary team professionals' perspective about adherence during the COVID-19 pandemic. **Methods:** this is a cross-sectional study, carried out from December 2020 to July 2021, with professionals from the Intensive Care Unit of a university hospital. Bivariate and multivariate analyzes were performed to assess factors associated with professionals' knowledge. **Results:** of the 61 professionals interviewed, 85.2% were nursing professionals and 14.7% were physical therapists. Physical therapists obtained a higher percentage of correct answers in relation to the VAP prevention bundle items. Nursing professionals have little knowledge about the elements that make up the bundle, highlighting the need for training throughout the entire process of training and professional performance. Lack of training was the main difficulty reported (14.75%). Age was a factor associated with knowledge of the bundle ( $p=0.017$ ). **Conclusion:** there is a lack of knowledge and adherence to the VAP prevention bundle among nursing professionals. However, these professionals demonstrated a willingness to receive training to improve bundle adherence rates and current good VAP prevention practices.

**Keywords:** Pneumonia, Ventilator-Associated. Intensive Care Units. Pandemics. Infection Control. COVID-19.

### RESUMO

**Justificativa e Objetivos:** a COVID-19 aumentou a demanda pela terapia intensiva e intensificou a necessidade de ventilação mecânica e a gravidade da pneumonia associada à ventilação mecânica (PAV). A adoção de medidas

preventivas da PAV no contexto da pandemia tem grande impacto para salvar vidas. Assim, objetivou-se avaliar o conhecimento referido sobre o *bundle* de prevenção da PAV e seus fatores associados e a percepção dos profissionais da equipe multiprofissional sobre a adesão em vigência da pandemia de COVID-19. **Métodos:** trata-se de estudo transversal, realizado no período de dezembro de 2020 a julho de 2021, com profissionais da Unidade de Terapia Intensiva de hospital universitário. Foram realizadas análises bivariadas e multivariadas para avaliar os fatores associados ao conhecimento dos profissionais. **Resultados:** dos 61 profissionais entrevistados, 85,25% eram profissionais de enfermagem e 14,75% eram fisioterapeutas. Os fisioterapeutas obtiveram maior porcentagem de acertos em relação aos itens do *bundle* de prevenção da PAV. Os profissionais de enfermagem possuem pouco conhecimento sobre os elementos que compõem o *bundle*, remetendo à necessidade de capacitação ao longo de todo o processo de formação e atuação profissional. Falta de treinamento foi a principal dificuldade relatada (14,75%). A idade foi um fator associado ao conhecimento do *bundle* ( $p=0,017$ ). **Conclusão:** há deficiência no conhecimento e na adesão ao *bundle* de prevenção de PAV entre os profissionais de enfermagem. No entanto, esses profissionais demonstraram disposição em receber capacitação para melhoria das taxas de adesão ao *bundle* e das boas práticas de prevenção da PAV vigentes.

**Descritores:** *Pneumonia Associada à Ventilação Mecânica. Unidades de Terapia Intensiva. Pandemias. Controle de Infecções. COVID-19.*

## RESUMEN

**Justificación y Objetivos:** el cumplimiento de las medidas para prevenir la neumonía asociada a la ventilación mecánica (NAV) reduce significativamente esta infección hospitalaria, y el conocimiento de las medidas es fundamental para garantizar el cumplimiento. COVID-19 ha aumentado la demanda de cuidados intensivos y aumenta los riesgos y la gravedad de la NAV. Así, el objetivo fue evaluar los conocimientos referidos sobre el paquete de prevención de NAV y sus factores asociados y la percepción de los profesionales del equipo multidisciplinario sobre la adherencia durante la pandemia de COVID-19. **Métodos:** se trata de un estudio transversal, realizado de diciembre de 2020 a julio de 2021, con profesionales de la Unidad de Cuidados Intensivos de un hospital universitario. Se realizaron análisis bivariados y multivariados para evaluar los factores asociados al conocimiento de los profesionales. **Resultados:** de los 61 profesionales entrevistados, el 85,25% eran profesionales de enfermería y el 14,75% eran fisioterapeutas. Los fisioterapeutas obtuvieron un mayor porcentaje de respuestas correctas en relación con los ítems del paquete de prevención de NAV. Los profesionales de enfermería tienen poco conocimiento sobre los elementos que componen el paquete, destacándose la necesidad de capacitación durante todo el proceso de formación y desempeño profesional. La falta de formación fue la principal dificultad reportada (14,75%). La edad fue un factor asociado al conocimiento del paquete ( $p=0,017$ ). **Conclusión:** existe falta de conocimiento y adherencia al paquete de prevención de NAV entre los profesionales de enfermería. Sin embargo, estos profesionales demostraron voluntad de recibir capacitación para mejorar las tasas de cumplimiento del paquete y las buenas prácticas actuales de prevención de NAV.

**Palabras Clave:** *Neumonía Asociada al Ventilador. Unidades de Cuidados Intensivos. Pandemias. Control de Infecciones. COVID-19.*

## INTRODUCTION

The clinical picture of patients with COVID-19 can range from mild flu-like symptoms to more severe conditions, with systemic manifestations.<sup>1</sup> One of the most frequent complications of severe COVID-19 is acute respiratory distress syndrome (ARDS), causing diffuse alveolar damage, pulmonary capillary thrombi, and endothelial dysfunction, culminating in lung collapse and respiratory failure. This syndrome is the main indication for invasive mechanical ventilation (IMV) in COVID-19.<sup>2</sup>

When subjected to IMV for more than 48 hours, patients are at risk of acquiring ventilator-associated pneumonia (VAP), which occurs when there is contamination of the lungs by exogenous or endogenous bacteria.<sup>3</sup> The occurrence of VAP prolongs the time of ventilator use, increases the length of hospital stay and hospitalization costs, and can contribute to a higher risk of death.<sup>3</sup> Therefore, preventive measures for VAP, offered through a

bundle, are essential for the care of critically ill patients with COVID-19 on mechanical ventilation.

The VAP prevention bundle is defined as a set of evidence-based interventions that, when implemented in combination, reduce the risk of VAP.<sup>4</sup> The implementation of the VAP prevention bundle has resulted in a significant reduction in VAP incidence worldwide. In a hospital in Turkey, the VAP rate decreased from 15.91 per 1,000 ventilator days to 8.50 per 1,000 ventilator days.<sup>5</sup> In a hospital in Taiwan, VAP incidence density decreased significantly from 3.3 to 1.4 cases per 1,000 ventilator days after the bundle implementation.<sup>6</sup>

Likewise in Brazil, studies have shown that the bundle implementation reduces the number of VAP cases, mortality and hospital costs, improving the quality of care for patients in intensive care.<sup>7-8</sup>

The increase in new cases of severe pneumonia caused by COVID-19 has led to overcrowding and pressure

in Intensive Care Units (ICUs), resulting in a high rate of mechanical ventilation use. Moreover, it has changed the way these units operate, requiring new isolation spaces, new facilities and urgent hiring to meet this demand.<sup>9</sup>

The increased need for IMV leads to an increased risk of VAP, which, for patients with COVID-19 and ARDS, increases the severity of patients' condition and the chance of shock, septicemia, and death.<sup>10</sup> Therefore, the need to apply the VAP prevention bundle becomes even more expressive to reduce complications and promote patient recovery.

Thus, during the COVID-19 pandemic, in which ICU utilization rates reached 100% of their capacity, it is important to investigate the healthcare professionals' knowledge about the VAP prevention bundle and their adherence in this context. Identifying barriers to adherence can help establish measures to strengthen the application of these care measures, thus helping to reduce VAP rates in ICUs. Thus, the objective of this study was to assess the knowledge reported on the VAP prevention bundle and its associated factors and the perception of nursing and physical therapy team professionals regarding adherence, during the COVID-19 pandemic, in a hospital in Goiânia, Goiás.

## METHODS

### Study design, period and site

This is a cross-sectional study conducted in an ICU specifically for the care of patients with COVID-19 in a large hospital in Goiânia, Goiás, Brazil, from January to December 2021.

### Sample and eligibility criteria

The study population consisted of professionals directly involved in the implementation of the actions that make up the VAP prevention bundle, such as nurses, nursing technicians and physical therapists. The sampling process for this study was by convenience, in which all ICU professionals were invited to participate, following the following inclusion criteria: having worked for more than five months in the unit and being part of the nursing or physical therapy team. Participants who responded less than 50% of the instrument were excluded. Of the 80 professionals eligible for the study, 71 agreed to participate; of these, ten were excluded due to incomplete completion of the instrument. Thus, the sample consisted of 61 professionals, representing 76.2% of the intended population.

### Data collection

Data were collected from January to August 2021. Professionals were contacted personally at their work unit and invited to participate in the study. Those who accepted agreed to participate with the data collection team, depending on their availability to respond to the instrument. Data were collected in a room attached to the unit, where the professional responded to the instrument and, at the end, returned it to the researcher.

The data collection instrument was structured in

three parts, namely: 1. Sociodemographic data; 2. Professional participation in continuing education activities addressing VAP prevention measures and their perception of adherence to the bundle during the pandemic (dichotomous closed-ended questions); 3. Open-ended question: "What interventions make up the VAP prevention bundle?". The instrument was developed by the research team, based on the Brazilian National Health Regulatory Agency (In Portuguese, *Agência Nacional de Vigilância Sanitária* - ANVISA) Healthcare-Associated Infection Prevention Measures.<sup>11-12</sup> It was submitted for analysis by three judges with PhDs in nursing and working in healthcare-associated infection control (HAI) services. After adjustments, the instrument was validated by a pilot test in the ICU of another institution.

### Study variables

The dependent variable was knowledge of elements that make up the VAP prevention bundle (oral hygiene, elevated head of the bed, checking cuff pressure, possibility of extubation, daily awakening/reduction of sedation and adequate ventilation system maintenance). For each participant, the number of items that make up the bundle was counted, categorized as: knows up to three elements of the bundle and knows more than three elements of the bundle. Independent variables were divided into sociodemographic and professional data, and information related to participation in continuing education activities, detailed in Table 1.

### Data analysis

The data were analyzed using the statistical program R version 4.3.1. Descriptive analyses were performed with presentation of qualitative data in the form of absolute and relative frequencies and quantitative data in the form of median and interquartile range. The relative frequencies with respective confidence intervals on the incentive to adhere to the prevention bundle and knowledge of each element were presented in the form of a bar graph. The chi-square test for trend was used to compare the encouragement to adhere to the bundle before the pandemic with the period during the pandemic. Logistic regression models were used to perform bivariate and multivariate analyses of factors associated with knowledge of elements of the VAP prevention bundle, from which unadjusted and adjusted Odds Ratios (OR) were obtained.

Variables that presented  $p < 0.2$  were included in the multivariate model, so the stepwise strategy was established as the input method for these variables into the model, in which they are included in the model so that the adjustment identifies whether it is feasible to include or remove them, with the model starting without any variables until a model with a better fit is identified. As a criterion for defining the model's fit, we established the Akaike Information Criterion (AIC). Thus, the final model included only the variables that fit best. Variables that presented a  $p$ -value  $< 0.05$  were considered statistically significant.


The study protocol was submitted to and approved by the Research Ethics Committee of the *Hospital*



**Table 2.** Sociodemographic characteristics and adherence to the ventilator-associated pneumonia prevention bundle among professionals in an Intensive Care Unit for COVID-19 patients participating in the study. Goiânia, GO, Brazil, 2021.



| Variables   | N(%)         |
|---|--------------|
| <b>Sex</b>  |              |
| Female  | 52 (85.2%)   |
| Male  | 9 (14.8%)    |
| <b>Profession</b>   |              |
| Nursing technicians   | 31 (50.8%)   |
| Nurses  | 21 (34.4%)   |
| Physical therapists   | 9 (14.8%)    |
| <b>Other employment relationship</b>                                  |              |
| Yes   | 48 (78.7%)   |
| No  | 13 (21.3%)   |
| <b>Participation in training on the bundle (in the unit)</b>          |              |
| Yes   | 19 (31.1%)   |
| No  | 42 (68.9%)   |
| <b>Participation in training on the bundle (in the other link)</b>    |              |
| Yes   | 14 (22.9%)   |
| No  | 47 (77.1%)   |
| <b>Felt encouraged to apply the bundle (before the pandemic)</b>      |              |
| Yes   | 40 (65.6%)   |
| No  | 21 (34.4%)   |
| <b>Felt encouraged to apply the bundle (during the pandemic)</b>      |              |
| Yes   | 57 (93.4%)   |
| No  | 4 (6.6%)     |
| <b>Believe that the pandemic influenced the bundle implementation</b> |              |
| Yes   | 40 (65.4%)   |
| No  | 21 (34.4%)   |
| <b>Believe that the pandemic influenced the bundle completion</b>     |              |
| Yes   | 39 (65%)     |
| No  | 21(35%)      |
| <b>Age (years) [median (IQR*)]</b>                                    |              |
|   | 50 (47 – 54) |
| <b>Time since graduation [median (IQR*)]</b>                          |              |
|   | 19 (14 – 25) |
| <b>Job tenure in the unit [median (IQR*)]</b>                         |              |
|   | 12 (6 – 15)  |

Note: \*IQR = interquartile range.



Note: \*Chi-square test for trend.

**Figure 1.** The COVID-19 pandemic increased the frequency of professionals who felt encouraged to adhere to the ventilator-associated pneumonia prevention bundle. Goiânia, GO, Brazil, 2020-2021.



Note: A: nursing professionals; B: physical therapists; E1: oral hygiene; E2: elevated head of bed; E3: checking cuff pressure; E4: possibility of extubation; E5: daily awakening/reduction of sedation; E6: adequate ventilation system maintenance.

**Figure 2.** Knowledge of ventilator-associated pneumonia prevention bundle elements among Intensive Care Unit professionals during the COVID-19 pandemic. Goiânia, GO, Brazil, 2021.

th age, participation in training on bundle V in the other employment relationship, feeling encouraged to apply the bundle and believing that the pandemic influenced the bundle implementation. In the multivariate analysis, only age remained statistically significant, showing that the younger the age, the greater the knowledge about the bundle (OR: 0.83; 95% CI: 0.71 – 0.96; p=0.017) (Table 3).

## DISCUSSION

VAP is one of the most common HAIs in ICUs, and can worsen patients' clinical conditions and increase their chances of death. However, correctly implemented prevention measures can reduce the incidence of this infection and its impacts.<sup>6</sup>

The COVID-19 pandemic, in which the management of the severe form of the disease requires ventilatory support in the ICU, has caused overcrowding, increased rates of mechanical ventilation use and contributed to the increased risk of VAP incidence.<sup>13</sup> Therefore, it is essential to diagnose the knowledge and practice of professionals

**Table 3.** Factors related to knowledge of ventilator-associated pneumonia prevention bundle elements among Intensive Care Unit professionals during the COVID-19 pandemic. Goiânia, GO, Brazil, 2020-2021.

| Variables  | Knowledge [n (%)]         |                                | Unadjusted OR*<br>(95% CI) | p-value | Adjusted OR*<br>(95% CI) | β†     | p-value‡ |
|--|---------------------------|--------------------------------|----------------------------|---------|--------------------------|--------|----------|
|  | > three elements<br>(n=6) | Up to three<br>elements (n=55) |                            |         |                          |        |          |
| <b>Sex</b>   |                           |                                |                            |         |                          |        |          |
| Female   | 5 (9.6)                   | 47 (90.4)                      | 0.85 (0.08; 8.27)          | 0.889   |                          |        |          |
| Male   | 1 (11.1)                  | 8 (88.9)                       | 1                          |         |                          |        |          |
| <b>Age [median (IQR§)]</b>   | 36.5 (33.2; 43.5)         | 51.0 (49.0; 54.0)              | 0.80 (0.69; 0.93)          | 0.003   | 0.83 (0.71; 0.96)        | -0.185 | 0,017    |
| <b>Profession</b>  |                           |                                |                            |         |                          |        |          |
| Nurse  | 1 (4.8)                   | 20 (95.2)                      | -                          | 0.998   |                          |        |          |
| Physical therapist   | 5 (55.6)                  | 4 (44.4)                       | -                          | 0.997   |                          |        |          |
| Nursing technician   | 0 (0.0)                   | 31 (100.0)                     | 1                          |         |                          |        |          |
| <b>Another employment relationship</b>   |                           |                                |                            |         |                          |        |          |
| Yes  | 6 (12.5)                  | 42 (87.5)                      | -                          | 0.995   |                          |        |          |
| No   | 0 (0.0)                   | 13 (100.0)                     | 1                          |         |                          |        |          |
| <b>Time since training [median (IQR§)]</b>   | 15.0 (10.7; 17.7)         | 6.8 (14.0; 25.0)               | 0.84 (0.71; 1.00)          | 0.052   |                          |        |          |
| <b>Job tenure in the unit [median (IQR§)]</b>                                      | 7.5 (6.2; 13.2)           | 12.0 (6.0; 15.0)               | 0.94 (0.79; 1.12)          | 0.531   |                          |        |          |
| <b>Participation in training on the bundle (in the unit) [during the pandemic]</b> |                           |                                |                            |         |                          |        |          |
| Yes  | 1 (5.26)                  | 18 (94.7)                      | 0.41 (0.04; 3.78)          | 0.433   |                          |        |          |
| No   | 5 (11.9)                  | 37 (88.1)                      | 1                          |         |                          |        |          |
| <b>Participation in training on the bundle (in the other link)</b>                 |                           |                                |                            |         |                          |        |          |
| Yes  | 4 (28.6)                  | 10 (71.4)                      | 8.99 (1.44; 56.12)         | 0.018   |                          |        |          |
| No   | 4 (4.3)                   | 45 (95.7)                      | 1                          |         |                          |        |          |
| <b>Feel encouraged to apply the bundle</b>   |                           |                                |                            |         |                          |        |          |
| Yes  | 4 (7.0)                   | 53 (93.0)                      | 0.07 (0.01; 0.68)          | 0.021   |                          |        |          |
| No   | 2 (50.0)                  | 2 (50.0)                       | 1                          |         |                          |        |          |
| <b>Believe that the pandemic influenced the bundle implementation</b>              |                           |                                |                            |         |                          |        |          |
| Yes  | 1 (2.5)                   | 39 (97.5)                      | 0.08 (0.01; 0.75)          | 0.027   | 0.17 (0.01; 1.89)        | -1.76  | 0,149    |
| No   | 5 (23.8)                  | 16 (76.2)                      | 1                          |         | 1                        |        |          |
| <b>Believe that the pandemic influenced the bundle completion</b>                  |                           |                                |                            |         |                          |        |          |
| Yes  | 2 (5.1)                   | 37 (94.9)                      | 0.23 (0.04; 1.37)          | 0.107   |                          |        |          |
| No   | 4 (19.1)                  | 17 (80.9)                      | 1                          |         |                          |        |          |

Note: \*OR = Odds Ratio; †β = model coefficients; ‡p-value = significance level; §IQR = interquartile range; model fit parameter: AIC=53.89.

in the application of VAP prevention measures which, if well targeted, can significantly reduce this complication.<sup>6,14</sup>

In the ICU assessed in this study, the physical therapy team was directly responsible for registering the bundle, but most studies regarding adherence to this tool, in different parts of the world, are aimed at the nursing team,<sup>15-16</sup> which highlights the leading role of this professional team in the application of VAP preventive measures. Thus, our study innovated by including physical therapists as participants, in addition to nursing professionals, due to their institutional responsibility in the bundle application and registration and their recognized interprofessional performance in VAP prevention.<sup>17</sup>

Only 26% of professionals reported having received training on the VAP prevention bundle at the institution. Since VAP is one of the main adverse events for critical care patients, it would be essential for workers in this sector to be periodically updated. The number of professionals who received training at other institutions was also low,

showing that not only at the study institution, but also in other services, investment in continuing education for VAP prevention is insufficient.

It has been shown that patients with COVID-19 are at higher risk of developing VAP<sup>21</sup> and that VAP in patients with COVID-19 was associated with greater complications, such as shock (71% vs. 48%, p = 0.009), ICU death (52% vs. 30%, p = 0.011), and bacteremia (26% vs. 13%, p = 0.038), than patients who develop VAP without COVID-19.<sup>10</sup> These findings make the application of VAP prevention bundles essential in the context of the COVID-19 pandemic.

Considering the importance of applying VAP prevention bundles, some factors can be suggested for the low number of professionals who reported training in VAP prevention, such as a shortage of professionals, due to the expansion in the number of beds, and high turnover, due to the replacement of infected professionals, generating emergency hiring of professionals with little

experience, in addition to a large number of new management and assistance demands related to preventive measures against infection in the unit.

Before the pandemic, only 65% of survey participants reported feeling encouraged to apply the bundle, a worrying fact, since this infection is preventable and has harmful consequences for patients and healthcare services, increasing their costs.<sup>8-11</sup>

It is noteworthy that, during the pandemic, most professionals felt more encouraged to apply the VAP prevention bundle (93%), compared to the period before the pandemic ( $p < 0.001$ ). It is important to highlight that the variable feeling encouraged to apply the bundle was a self-report by participants and does not reflect real adherence, which was not the target of this investigation. Despite this, low knowledge among nursing professionals was evidenced, which supports the lack of training reported in the interviews, contrasting with the increased encouragement to apply the bundle during the pandemic. In practice, this may represent inconsistent care and reinforces the need for training teams, since feeling encouraged to apply it may result in greater interest in learning. However, this was a period marked, at times, by greater adherence to protective measures, resulting in a reduction in healthcare-associated infections, and, at times, by relaxation and an increase in HAI rates.<sup>19,20</sup>

The practices selected for knowledge assessment correspond to six care procedures that make up the bundle: 1 - positioning of the headboard between 30° and 45°; 2 - checking the cuff pressure between 20-30 cmH<sub>2</sub>O; 3 - oral hygiene with chlorhexidine 0.12%; 4 - respiratory and motor physical therapy; 5 - care with the aspiration of secretions; 6 - interruption of sedative infusion.<sup>11,21</sup>

The study showed that nursing professionals at the institution have little knowledge about the elements that make up the VAP prevention bundle, which certainly limits their ability to implement prevention actions. Some of the components of the bundle are routine nursing care, such as oral hygiene and positioning. However, it is necessary to understand these actions as part of the bundle of VAP preventive measures. A study conducted in Australia,<sup>16</sup> which assessed the knowledge of specialist nurses in ICUs, identified that the most mentioned element of the bundle for VAP prevention was positioning (90.9), unlike our study, in which this element was mentioned by only 28% of professionals, referred to here as elevated headboard, a highly effective nursing intervention for VAP prevention.<sup>22</sup>

Elevated headboard position, oral hygiene and adequate maintenance of cuff pressure represent the main preventive measures for subglottic aspiration, one of the most common causes of VAP.<sup>4,11</sup> They also represent routine nursing interventions and, even so, were rarely mentioned as measures to prevent VAP. Furthermore, the assessment of patients, which involves the level of consciousness and oxygenation, is also part of nursing routine, which, together, covers the other items in the bundle, such as the possibility of extubation, awakening/reduction of sedation and adequate ventilation system

maintenance. However, despite being nursing interventions, they were not mentioned as components of the VAP prevention bundle.

Considering all the elements of the bundle, in general, studies have shown a low level of knowledge among professionals, which certainly impacts adherence to the VAP bundle and incidence.<sup>15,16</sup> A study conducted in Brazil found that the frequency of adherence to the bundle by nursing professionals was significant only for three proposed items, namely the position of the headboard between 30° and 45°, care with suction of secretions and respiratory and motor physical therapy, with the overall adherence rate being 77.4%; therefore, no events of complete bundle implementation were observed.<sup>23</sup> Such data support the results of this study, and alert ICU managers to team continued education.

A study conducted in Porto Alegre (RS) assessed nursing adherence to the VAP prevention bundle and the incidence rate before and after implementing a continuing education program. The data revealed an increase in adherence to all components of the interventions proposed in the bundle and, consequently, a reduction in VAP incidence rates, which demonstrates that continuing education is an important tool for maintaining adherence to the VAP prevention bundle in favor of the safety of critically ill patients in the ICU.<sup>8</sup> It is worth noting that, in the present study, physical therapists were more prepared, when compared to the nursing team, in relation to all elements of the VAP prevention bundle. It is also worth noting that care should be provided in a multidisciplinary manner, resulting in substantial improvements in healthcare.<sup>17</sup>

Logistic regression showed that, in the present study, age was inversely related to knowledge of bundle elements, suggesting that younger professionals tend to have more knowledge; this may be due to the recent inclusion of topics related to patient safety, especially in HAI prevention, in healthcare professional training.<sup>24-25</sup> Therefore, it is essential to recognize the importance of HAI prevention and control actions in professional training, ensuring this content in the training curriculum for healthcare professionals, as well as prioritizing professional training, providing instruments for monitoring adherence to the bundle, in order to reduce the incidence of VAP which, especially for patients with COVID-19, culminates in unfavorable outcomes.

This study has limitations, such as the distribution of participants from the physical therapy team being smaller than that of the nursing team, but it represents the proportion between the professional categories. Self-reported knowledge may also contribute to limitations of this study, as it may not reflect real knowledge on the subject, indicating the need for further research with more objective assessments of knowledge.

Although this study was conducted in the context of the COVID-19 pandemic, the results have the potential to qualify ICU care in different contexts, and are especially transferable to other infectious or non-infectious respiratory impairment syndromes that may increase the demand for ICUs, signaling the need for training in

the face of new hires, and the maintenance of a qualified team to apply VAP prevention measures.

## REFERENCES

1. Li L, quan, Huang T, Wang Y, qing, et al. COVID-19 patients' clinical characteristics, discharge rate, and fatality rate of meta-analysis. *J Med Virol* 2020; 1;92(6):577–83. <https://doi.org/10.1002/jmv.25757>
2. Attaway AH, Scheraga RG, Bhimraj A, et al. Severe covid-19 pneumonia: pathogenesis and clinical management. *BMJ* 2021; 10;372. <https://doi.org/10.1136/bmj.n436>
3. Papazian L, Klompas M, Luyt CE. Ventilator-associated pneumonia in adults: a narrative review. *Intensive Care Med*. 2020; 46(5):888–906. <https://doi.org/10.1007/s00134-020-05980-0>
4. Wip C, Napolitano L. Bundles to prevent ventilator-associated pneumonia: how valuable are they? *Curr Opin Infect Dis*. 2009;22(2). <https://doi.org/10.1097/qco.0b013e3283295e7b>
5. Okgün Alcan A, Demir Korkmaz F, Uyar M. Prevention of ventilator-associated pneumonia: Use of t care bundle approach. *Am J Infect Control*. 2016;44(10):e173–6. <https://doi.org/10.1016/j.ajic.2016.04.237>
6. Lim KP, Kuo SW, Ko WJ, et al. Efficacy of ventilator-associated pneumonia care bundle for prevention of ventilator-associated pneumonia in the surgical intensive care units of a medical center. *J Microbiol Immunol Infect* 2015;48(3):316–21. <https://doi.org/10.1016/j.jmii.2013.09.007>
7. Ferreira CR, de Souza DF, Cunha TM, et al. The effectiveness of a bundle in the prevention of ventilator-associated pneumonia. *Brazilian J Infect Dis* 2016; 20:267–71. <https://doi.org/10.1016/j.bjid.2016.03.004>
8. Branco A, Lourençone EMS, Monteiro AB, Fonseca JP, Blatt CR, Caregnato RCA. Education to prevent ventilator-associated pneumonia in intensive care unit. *Rev Bras Enferm* 2020;73. <https://doi.org/10.1590/0034-7167-2019-0477>
9. Dhala A, Gotur D, Hsu SHL, et al. A Year of Critical Care: The Changing Face of the ICU During COVID-19. *Methodist Debakey Cardiovasc J* 2021;17(5):31. <https://doi.org/10.14797/mdcvj.1041>
10. Rouyer M, Strazzulla A, Youbong T, et al. Ventilator-associated pneumonia in COVID-19 patients: A retrospective cohort study. *Antibiotics* 2021;10(8):988. <https://doi.org/10.3390/antibiotics10080988>
11. Brasil. Agência Nacional de Vigilância Sanitária. Medidas de prevenção de infecção relacionada à assistência à saúde [Internet]. Brasília; 2017. <https://www.gov.br/anvisa/pt-br/centraisdeconteudo/publicacoes/servicosdesaude/publicacoes/caderno-4-medidas-de-prevencao-de-infeccao-relacionada-a-assistencia-a-saude.pdf/view>
12. Brasil. Agência Nacional de Vigilância Sanitária. Nota Técnica GVIMS/GGTES/ANVISA, n. 04/2020. Orientações para serviços de saúde: medidas de prevenção e controle que devem ser adotadas durante a assistência aos casos suspeitos ou confirmados de infecção pelo novo coronavírus (SARS-CoV-2) [Internet]. Brasília; 2021. [https://www.gov.br/anvisa/pt-br/centraisdeconteudo/publicacoes/servicosdesaude/notas-tecnicas/2020/nota-tecnica-gvims\\_ggtes\\_anvisa-04\\_2020-25-02-para-o-site.pdf](https://www.gov.br/anvisa/pt-br/centraisdeconteudo/publicacoes/servicosdesaude/notas-tecnicas/2020/nota-tecnica-gvims_ggtes_anvisa-04_2020-25-02-para-o-site.pdf)
13. Wicky PH, Niedermann MS, Timsit JF. Ventilator-associated pneumonia in the era of COVID-19 pandemic: How common and what is the impact? *Crit Care* 2021;25(1):1–3. <https://doi.org/10.1186/s13054-021-03571-z>
14. Vazquez Guillamet C, Kollef MH. Is Zero Ventilator-Associated Pneumonia Achievable?: Practical Approaches to Ventilator-Associated Pneumonia Prevention. *Clin Chest Med* 2018;39(4):809–22. <https://doi.org/10.1016/j.ccm.2018.08.004>
15. Dehghan M, Arab M, Akafzadeh T, et al. Intensive care unit registered nurses' perceived barriers towards ventilated associated pneumonia prevention in southeast Iran: a cross-sectional descriptive – an analytical study. *BMJ Open* 2022 Sep 1;12(9):e064147. <https://doi.org/10.1136/bmjopen-2022-064147>
16. Madhuvu A, Endacott R, Plummer V, et al. Nurses' knowledge, experience and self-reported adherence to evidence-based guidelines for prevention of ventilator-associated events: A national online survey. *Intensive Crit Care Nurs* 2020;59:102827. <https://doi.org/10.1016/j.iccn.2020.102827>
17. Ntinga MN, van Aswegen H. Physiotherapists' perceptions of collaborations with inter-professional team members in an ICU setting. *South African J Crit care Off J Crit Care Soc* 2020;36(2). <https://doi.org/10.7196/sajcc.2020.v36i2.431>
18. Maes M, Higginson E, Pereira-Dias J, et al. Ventilator-associated pneumonia in critically ill patients with COVID-19. *Crit Care* 2021;25(1):25. <https://doi.org/10.1186/s13054-021-03460-5>
19. Arabi YM, Azoulay E, Al-Dorzi HM, Phua J, Salluh J, Binnie A, et al. How the COVID-19 pandemic will change the future of critical care. *Intensive Care Med* 2021;47(3):282–91. <https://doi.org/10.1007/s00134-021-06352-y>
20. O'Toole RF. The interface between COVID-19 and bacterial healthcare-associated infections. *Clin Microbiol Infect* 2021;27(12):1772–6. <https://doi.org/10.1016/j.cmi.2021.06.001>
21. Kalil AC, Metersky ML, Klompas M, et al. Management of Adults With Hospital-acquired and Ventilator-associated Pneumonia: 2016 Clinical Practice Guidelines by the Infectious Diseases Society of America and the American Thoracic Society. *Clin Infect Dis* 2016 Sep 1;63(5):e61–111. <https://doi.org/10.1093/cid/ciw353>
22. Güner CK, Kutlutürkan S. Role of head-of-bed elevation in preventing ventilator-associated pneumonia bed elevation and pneumonia. *Nurs Crit Care* 2022 Sep 1;27(5):635–45. <https://doi.org/10.1111/nicc.12633>
23. Barros FRB de. Adesão ao bundle de prevenção de pneumonia associada à ventilação mecânica. *Rev Cuid* 2019;10(2). <https://doi.org/10.15649/cuidarte.v10i2.746>
24. Ribeiro G, Pires DEP de, Martins MM, et al. Biosafety and patient safety: the perspective of nursing teachers and students. *Acta Paul Enferm* 2023;36. <https://doi.org/10.37689/actaape/2023AO029211>
25. Wegner W, Silva SC da, Kantorski KJC, et al. Education for culture of patient safety: Implications to professional training. *Esc Anna Nery* 2016;20. <https://doi.org/10.5935/1414-8145.20160068>



## AUTHORS' CONTRIBUTIONS

**Hélio Galdino-Júnior** contributed to the conception, study design and project administration, bibliographic research, interpretation and description of results, manuscript writing and article critical review. **George Oliveira** contributed to the interpretation and description of results, statistical analysis, manuscript writing and final review. **Ana Clara Alves Campos** contributed to

bibliographic research, field data collection, abstract writing, methodology, interpretation of results, conclusions. **Jéssica de Oliveira Montebello** contributed to bibliographic research, field data collection, abstract writing, methodology, interpretation of results, conclusions.

All authors have approved the final version to be published and are responsible for all aspects of the work, including ensuring its accuracy and integrity.